



E-ISSN: 2663-1067
P-ISSN: 2663-1075
IJHFS 2022; 4(2): 96-100
Received: 11-04-2022
Accepted: 16-05-2022

Olamide Falaiye
Center of Excellence in
Agricultural Development and
Sustainable Environment,
Federal University of
Agriculture, Abeokuta, Ogun
State, Nigeria

Jacob Goke Bodunde
Department of Horticulture,
Federal University of
Agriculture, Abeokuta, Ogun
State, Nigeria

Eyitayo Adekunle Makinde
Department of Horticulture,
Federal University of
Agriculture, Abeokuta, Ogun
State, Nigeria

Olajide Philip Sobukola
Department of Food Science
and Technology, Federal
University of Agriculture,
Abeokuta, Ogun State, Nigeria

Asimiu Kasali
Center of Excellence in
Agricultural Development and
Sustainable Environment,
Federal University of
Agriculture, Abeokuta, Ogun
State, Nigeria

Oluwatosin Mary Peters
Department of Plant
Physiology and Crop
Production, Federal
University of Agriculture,
Abeokuta, Ogun State, Nigeria

Corresponding Author:
Olamide Falaiye
Center of Excellence in
Agricultural Development and
Sustainable Environment,
Federal University of
Agriculture, Abeokuta, Ogun
State, Nigeria

The effect of spacing on the growth, phenology and yield of two cucumber (*Cucumis sativus* L.) varieties in the forest-savanna: Transition agro-ecological zone of Nigeria

Olamide Falaiye, Jacob Goke Bodunde, Eyitayo Adekunle Makinde, Olajide Philip Sobukola, Asimiu Kasali and Oluwatosin Mary Peters

Abstract

The increasing population in Nigeria has resulted in a high demand for food. Some of cucumber production constraints include limited high yielding varieties and appropriate spacing. Therefore, two field trials were conducted to determine growth and yield of two cucumber varieties in 2019, at the research farm of Federal University of Agriculture Abeokuta (Latitude 7°15'N and Longitude 3°25'E). The experimental design was a three-time replicated Randomized Complete Block Design (RCBD) in a split-plot layout. The main and the sub plots were given the factors: variety and spacing, respectively. The data was analyzed using ANOVA, and the means of significant treatments were separated using Least Significant Difference ($p < 0.05$). Monalisa produced significantly ($p < 0.05$) longer vines than CU 999 in the early season. The results of the experiment showed that increase in plant density brought about increase in fruit yield.

Keywords: Cucumber, variety, spacing, growth, yield

1. Introduction

Cucumber (*Cucumis sativus* L.) is one of Nigeria's most popular fruits and vegetables. It is the world's fourth most farmed vegetable and is often regarded as one of the healthiest foods available [1]. It belongs to the Cucurbitaceae family and is one of the most popular members. Cucumber is a South Asian native that is now grown on practically every continent. Cucumbers are available in a wide range of forms and sizes and are sold all over the world [2, 3]. They are vine crops that are cultivated on the ground, on poles, or on anchored trellises to suspend fruit [2, 3]. Cucumber fruit is high in vitamins A, C, K, B6, potassium, dietary fiber, pantothenic acid, magnesium, and phosphorus [4]. Cucumber cultivation is growing increasingly popular in a large area of Nigeria, according to [5], possibly due to its strong nutritional and medicinal benefits, as well as its use as a component ingredient in pharmaceuticals [6].

Plant spacing is one of the most essential elements in crop production, according to [7], since proper crop spacing makes optimal use of resources by limiting competition among plants with similar cultural requirements. [8] Investigated the influence of plant spacing on cucumber yield in a protected environment and found that a plant spacing of 60 cm 60 cm produced the maximum yield when compared to 60 cm 30 cm or 90 cm 60 cm spacing. According to [9], increasing plant density from 2 to 10 plants per m² increased yield per plant but decreased productivity per unit area, whereas decreasing plant density increased yield per unit area [10]. Tested the effect of plant population on cucumber yield and fruit quality, and discovered that plant population had significant impact on cucumber yield. He found that as plant density increased, reproductive yield (kg/plant) decreased. The highest plant density resulted in the largest fruit output.

[11] Investigated the influence of spatial arrangement on three cucumber types grown in a controlled environment, finding that close spacing resulted in a high total and marketable yield per unit area. Growth and yield of plant all rose dramatically as plant spacing increased, according to an experiment done by [12, 5].

Investigated the impact of plant spacing on cucumber growth and yield in a protected environment and found that the closest plant spacing (50 cm x 30 cm) produced the most fruits, and fruits with higher weight than the 50 cm x 40 cm spacing. This research was therefore done to determine the appropriate spacing for increased productivity to meet the consumers' demand.

2. Materials and Method

The experiment was carried out at the Directorate of University Farms' (DUFARMS) Teaching and Research site of the Federal University of Agriculture, Abeokuta, in the forest-savanna-transition agro-ecological zone. The rainfall distribution pattern for Abeokuta is bimodal, having the first mode between June and July and the second mode in September. The annual rainfall ranges from 1145 to 1270 mm. The experiment was conducted in two trials in 2019.

Soil samples collected from experimental sites were subjected to routine laboratory analysis before planting.

The treatments were arranged as a split plot fitted into a Randomized Complete Block Design with three replications using sub plot size of 2 m x 2 m. The treatment consisted of two cucumber varieties (CU 999 and Monalisa) and plant spacing (75 x 25, 75 x 50, 75 x 75) cm.

At 3, 4, and 5 WAS, primary vine length was measured from the soil surface to the tip of the stem of the five tagged plants in the center row using a meter rule. At 3, 4, and 4 WAS, the number of leaves on the sample plants were counted. Days to 50% flowering/blooming were calculated using the number of flowers observed on 50% of cucumber stands in each plots and days to fruiting were calculated using this same method. Weight of fruit was done using a top scale to weigh the fruits harvested from each net plot. Number of fruits harvested from the sample plants was counted at each harvest. Fruit girth was done using a veneer caliper while fruit length was done with a meter ruler.

3. Statistical analysis

The collected data were subjected to Analysis of Variance (ANOVA). At a 5% level of probability, the Least Significant Difference (LSD) was used to separate the means of significant treatments.

4. Results and Discussion

The experimental site's soil texture was sandy loam. The soil had a slightly acidic pH. The pH was (6.8) in both trials of 2019. The nitrogen content of the soil was medium (between 0.16% and 0.19%). Low nitrogen content is < 0.15%, medium is 0.15% to 0.20% and high is > 0.20%. The soil had a low organic matter content (Table 1).

Significant ($p \leq 0.05$) varietal difference existed in primary vine length of the cucumber varieties at 3, 4 and 5 WAS in the early season of 2019 while no difference was observed in the late season (Table 2). Monalisa variety had longer vines compared to CU 999. Spacing significantly ($p \leq 0.05$) influenced primary vine length at 3, 4 and WAS in the early season of 2019 and no such difference was observed in the late season. Plants in plots with spacing of 75 cm x 75 cm had longer vines.

Significant difference was observed on number of leaves of the varieties at 3, 4 and 5 WAS in the early season of 2019 (Table 3). In comparison to CU 999, the Monalisa variety produced plants with more leaves. At 3 and 4 WAS, spacing had a significant ($p \leq 0.05$) impact on the number of leaves in the early and late season while at 4 and 5 WAS, significant differences were observed only in the early season. Cucumber plants in plots with spacing of 75 cm x 75 cm

producing plants with higher number of leaves.

During the experiment, there were substantial differences in primary vine length and number of leaves/plant of cucumber; the disparities in growth rate indices are generally linked to their genetic make-up, according to [13]. This was in line with the findings of [14], who found that genetic variables improved plant height, leaf area, and pod output.

Variety had significant ($p \leq 0.05$) effect on days to 50% flowering and days to fruit set of cucumber in the early and late season of 2019 (Table 4). Variety CU 999 flowered earlier than Monalisa. Effect of spacing on days to 50% flowering and days to fruit set was significant ($p \leq 0.05$) in both seasons of 2019. Plants spaced at 75 cm x 75 cm flowered earlier than those spaced at 75 cm x 25 cm and 75 cm x 50 cm

Variety significantly ($p \leq 0.05$) influenced fruit girth in both seasons of 2019. CU 999 produced fruits with larger girth compared to Monalisa variety (Table 5). Effect of spacing on fruit girth was significant ($p \leq 0.05$) only in the late season. Plants spaced at 75 cm x 50 cm produced larger fruits while plants spaced at 75 cm x 75 cm had the smallest fruit girth. There was significant ($p \leq 0.05$) interaction between variety and spacing on fruit girth in both seasons.

Varietal influence was observed on fruit length in both seasons of 2019 (Table 5). Variety CU 999 produced significantly ($p \leq 0.05$) longer fruits than Monalisa. Fruit length was significantly ($p \leq 0.05$) influenced by spacing only in the late season. Plants spaced at 75 cm x 50 cm in the late produced longer fruits.

Variety significantly ($p \leq 0.05$) influenced fruit weight/plant in both seasons with CU 999 producing heavier cucumber fruits than Monalisa in both seasons (Table 5). Spacing significantly ($p \leq 0.05$) impacted fruit weight/plant in both seasons. Plant in plots spaced at 75 cm x 25 cm produced fruits with higher weight.

Variety had significant ($p \leq 0.05$) influence on number of fruits per plants in the two seasons with CU 999 producing more cucumber fruits than Monalisa in both seasons (Table 5). Spacing also had a significant ($p \leq 0.05$) impact in both seasons. Plant in plots spaced at 75 cm x 25 cm produced more fruits.

Varietal influence was observed on fruit yield (t/ha) in both seasons of 2019. Yield of CU 999 was higher compared to Monalisa (Table 5). Spacing significantly ($p \leq 0.05$) influenced yield (t/ha) in both seasons of 2019. In both seasons, plots spaced at 75 cm x 25 cm had higher yield.

Differential yield features were found in the CU 999 variety. Monalisa had a much lower number of fruits per plant, weight of fruits per plant, and total yield per hectare. Different cucumber researchers from around the world have reported on these differences in cucumber growth and production. The genetic composition of the types employed can be blamed for the discrepancies in vegetative and yield characteristics. The CU 999 type may have adapted to the surroundings more quickly than Monalisa. The CU 999 variety's vegetative features may have been more active, resulting in a robust source-to-sink interaction that led in the variety's high yields [15]. This was in line with [16] findings, which claimed that cucumber yield is influenced by genetic and environmental factors, and so varies depending on growing season and locale.

One of the most significant aspects of agricultural productivity is plant spacing. The highest plant density yielded the highest fruit production in this study. This opposed [17] findings, which showed that increasing plant spacing led to an increase in the no. of fruits/plant, fruit

length, and weigh of fruit/plant, while decreasing plant spacing led to increased plant height and no. of leaves. [18] Found that cassava ultimate leaf size and lateral shoot growth rose when planting density reduced, corroborating these findings. Plant density has a significant impact on growth and marketability of many fruit and vegetable crops, according to [19]. In their investigation of the influence of varied plant spacing on the output and quality of cucumbers in a greenhouse, [20] discovered that closer plant spacing resulted in a significantly lower fruit yield per plant. The increased fruit weight in closer plants could be attributable to greater assimilate use and increased assimilate allocation to the economic section. [21, 20, 22, 23, 24, 25, 26] all reported similar results. Plants with the widest spacing outperformed those that had smaller spacing in most of the growth indices studied. The higher fruit production could be attributed to the higher population density attained at a tight spacing of 75 cm x 25 cm, resulting in superior weed control through canopy shade, improved water usage

due to minimal rate of evapo-transpiration and better radiant energy utilization. According to [27], reduction in weed infestation, more oxygen supply and improved movement of water in the soil all contributed to increased okra development. The cucumber vines' failure to smother weeds due to a lack of plants and adequate room for weeds to thrive could explain the low yield achieved at wider spacing as crop and weed had to battle for available resources due to the vast quantity of space available. Plants in plots with spacing of 75 cm x 25 cm were determined to have the better yield, which was in line with [28], who discovered that the best spacing for optimum cucumber yield was 70 cm x 30 cm [29]. Also discovered that the closer the plant spacing, the higher the output [9]. Also found that when plant density grew from four to ten plants per meter square, the number of fruit declined. However, the results of the trial contradict those of [30], who found that higher plant spacing had the maximum yield.

Table 1: Physical and chemical properties of soil of the experimental sites

Properties	Early	Late
	Pre planting	Pre planting
pH	6.8	6.8
N (%)	0.18	0.19
Available P (mg/kg)	19.55	20.12
Org. C (%)	0.48	0.57
Org. M (%)	0.89	0.99
Ex. A (mEq/100g)	0.20	0.20
Na (cmol/kg)	0.28	0.32
k (cmol/kg)	0.40	0.43
Ca (cmol/kg)	0.30	0.32
Mg (cmol/kg)	0.38	0.39
Sand (%)	74.20	74.50
Clay (%)	7.20	6.10
Silt (%)	19.20	19.40
Textural Class	Sandy loam	Sandy loam

Table 2: Effect of Variety and Spacing on Primary Vine length of Cucumber at 3 - 5 Weeks after sowing

	3		4		5	
	Early	Late	Early	Late	Early	Late
Variety						
CU 999	28.12	23.31	176.43	75.94	357.63	128.27
Monalisa	32.23	24.14	222.25	66.11	440.81	115.45
LSD ($p \leq 0.05$)	1.118	NS	2.925	NS	14.001	NS
Spacing (cm)						
75 x 25	28.98	21.17	184.52	81.83	373.67	127.06
75 x 50	30.05	22.62	199.00	61.36	397.60	124.40
75 x 75	31.50	27.39	214.50	69.89	426.40	114.13
LSD ($p \leq 0.05$)	0.734	NS	1.920	NS	9.190	NS
V x S ($p \leq 0.05$)	NS	NS	NS	NS	NS	NS

Table 3: Effect of Variety and Spacing on Number of leaves of Cucumber at 3 - 5 Weeks after sowing

	3		4		5	
	Early	Late	Early	Late	Early	Late
Variety						
CU 999	11.66	8.42	21.57	15.06	39.36	19.76
Monalisa	13.48	10.76	24.90	12.71	47.79	15.61
LSD ($P \leq 0.05$)	0.182	NS	0.41	NS	0.615	NS
Spacing (cm)						
75 x 25	11.99	9.75	22.19	16.50	40.86	19.89
75 x 50	12.55	8.29	23.19	11.33	43.50	15.58
75 x 75	13.17	10.72	24.33	13.82	46.36	17.58
LSD ($p \leq 0.05$)	0.12	1.616	0.269	NS	0.403	NS
V x S ($p \leq 0.05$)	NS	NS	NS	NS	NS	NS

Table 4: Effect of Variety and spacing on phenology of cucumber

	Days to 50% flowering		Days to Fruit set	
	Early	Late	Early	Late
Variety				
CU 999	19.44	21.44	29.67	31.67
Monalisa	24.67	26.67	35.00	37.00
LSD ($p \leq 0.05$)	2.084	2.084	1.656	1.656
Spacing (cm)				
75 × 25	23.67	25.67	33.83	35.83
75 × 50	21.67	23.67	32.00	34.00
75 × 75	20.83	22.83	31.17	33.17
LSD ($p \leq 0.05$)	1.615	1.615	1.883	1.883
V × S ($p \leq 0.05$)	NS	NS	NS	NS

Table 5: Effect of Variety and spacing on fruit breadth, fruit length, Unit fruit weight, number of fruits and yield

	Fruit Girth		Fruit Length		Unit Fruit Weight		No of Fruits		Yield (t/ha)	
	Early	Late	Early	Late	Early	Late	Early	Late	Early	Late
Variety										
CU 999	56.27	56.55	24.94	25.92	3.85	5.91	10.28	19.25	9.63	14.79
Monalisa	47.32	52.39	19.19	22.79	1.30	1.93	5.39	7.35	3.25	4.81
LSD ($p \leq 0.05$)	1.153	2.055	4.235	1.550	1.596	1.971	2.426	5.879	3.99	4.929
Spacing (cm)										
75 × 25	51.63	50.07	21.89	24.29	3.52	7.14	10.92	20.47	8.80	17.85
75 × 50	52.37	57.31	22.15	26.46	3.11	2.26	9.00	9.45	7.78	5.64
75 × 75	51.39	56.03	22.16	22.31	1.10	2.36	3.58	9.98	2.75	5.91
LSD ($p \leq 0.05$)	NS	0.754	NS	0.448	1.574	0.533	3.907	1.318	3.935	1.332
V × S ($p \leq 0.05$)	4.856	0.907	NS	0.584	1.942	0.718	NS	2.018	4.854	1.794

5. Conclusions

From the study, it was established that variety CU 999 is higher yielding than Monalisa and the optimum growth and high fruit yield in cucumber is dependent on crop spacing in the field.

6. Acknowledgment

I'd want to thank the World Bank Group who sponsoring this research through the Federal University of Agriculture, Abeokuta, where one of the African Centre of Excellence in Agricultural Development and Sustainable Environment is located.

7. References

- Natural News. 10 Health benefits of cucumbers http://www.naturalnews.com/036769_cucumbers_health_benefits_rehydration.html. 2014.
- Nonneck IL. Vegetable production pub. Nan No. 1 Sotram Reinhold Company N.Y. 1989. p. 608-112.
- Wells J. Cucumber Business Leaves Family Farm in a Pickle. The Hamiton Spectator (Ontario, Canada) Retrieved 13 Nov, 2017; c2016.
- Olaniyi JO, Ogunbiyi EM, Alagbe DD. Effects of organo-mineral fertilizers on growth, yield and mineral nutrients uptake in cucumber. Journal of Animal & Plant Sciences. 2009;5(1):437-442.
- Nweke IA, Orji EC, Ijearu SI. The effect of staking and plant spacing on the growth and yield of cucumber (*Cucumis sativus* L.). J Environ Sci Tox Food Technol.(IOSR-JESTFT). 2013;3(4):26-31.
- Kumar D, Kumar S, Singh J, Rashmi N, Vashistha BD, Singh N. Free radical scavenging and analgesic activities of *Cucumis sativus* L. fruit extract. Journal of Young Pharmacology. 2010;2(4):365-368.
- Nnoke FN. Essentials of pedology and edaphology. Fedico Ventures, Abakaliki, Nigeria; c2001. p. 71.
- More TA, Chandra P, Majumdar G, Singh JK. Some observations on growing cucumber under plastic greenhouse. Proceedings of the 11th International Congress on the Use of Plastics in Agriculture, New Delhi, India; c1990 26th Feb-2nd Mar.
- Jacques LS, Jeronimo LA, Aron BH. Dry matter accumulation and partitioning of pickling cucumber plants under different plants densities. Clientia Rural Santa Maria. 2002;32:3-41.
- Nerson H. Plant density, fruit length and fruit type affect seed yield and quality in cucumber. Plant Density, Fruit Length and Fruit Type Affect Seed Yield and Quality in Cucumber; c2005. p. 1000-1007.
- Premalatha MGS, Wahundeniya KB, Weerakkody WAP, Wicramathunga CK. Plant training and spatial arrangement for yield improvements in greenhouse cucumber (*Cucumis sativus* L.) varieties; c2006.
- Lacob FC, Campeanu G, Atanasiu N. Effect of culture technique upon pickling cucumber hybrids in solarium tunnels in the Tartasesti- racan. Horticultura. 2009;53:133-136.
- Ibrahim R, Amans EB, Ahmed A, Abubakar IU. Growth and Yield of Tomato (*Lycopersicon esulentum* Mill) varieties influenced by crop spacing at Samaru, Northern Nigeria. Nig. J Hort. Sc. 2002;5:52-57.
- Sajjan AS, Shekhargounda M, Badanur. Influence of data of sowing, spacing and levels of nitrogen on yield attributes and seed yield of Okro. Ikamataka Journal of Agricultural Science. 2002;15(2):267-274.
- Cavatorra J, Moriarty G, Henning M, Glos M, Kreitingner M, Munger HM. A Monoecious Slicing Cucumber Inbred with Multiple Disease and Insect Resistances. Hort science. 2007;42(3):707-709.
- Staub JE, Bacher JJ. Cucumber as a processed vegetable (chapter six).Vegetable crops Research, USDA, University of Wisconsin Madison, WI; c2004.

- p. 129-193.
17. Serquen FC, Bacher J, Staub JE. Genetic analysis of yield components in cucumber at low plant density. *Journal of American Society of Horticulture Science*. 1997;122:522-528.
 18. Streck NA, Pinheiro DG, Zanon AJ, Gabriel LF, Rocha TS, Trevisan de Souza A and Rocha da Silva M. Effect of plant spacing on growth, development and yield of cassava in a subtropical environment. *Bragantia*; c2014, 73(4). Campinas Oct./Dec. [http:// dx.doi.org/10.1590/1678-4499.0159](http://dx.doi.org/10.1590/1678-4499.0159).
 19. Kosson R, Dobrzanska J. The effect of plant training system on yielding and storability of cucumber fruits. *Vegetable Crops Research Bulletin*. 2002;56:57-65.
 20. Echevarra PH, Castro AR. Influence of Different Plant Densities on the Yield and Quality of Greenhouse grown Cucumbers Grafted on Shintoza (*Cucurbita maxima* x *Cucurbita moschata*). *Acta Horticulturae*; c2002. p. 558. ed. Gaithersburg Maryland: AOAC International.
 21. Choudhari SM, More TA. Fertigation, fertilizer and spacing requirement of Tropical gynoeocious cucumber hybrids. *ISHS*. Tsukuba, Japan. *Acta Hort*. 2002;61:588.
 22. Motsenbocker CE, Arancibia RA. In-row spacing influences triploid watermelon yield and crop value. *Hort Technology*. 2002;12(3):437-440.
 23. Peil RM, Gonzalez-Real MM, Lopez-Galvez J. Light interception of a greenhouse cucumber crop: measurements and modelling results. *Acta horticulturae*; c2002.
 24. Khalid EA, Elwan MWM. Dependence of Pumpkin Yield on Plant Density and Variety. *American Journal of Plant Sciences*. 2011;2:636-643.
 25. Zhang F, Cui Z, Fan M, Zhang W, Chen X, Jiang R. Integrated soil crop system management: reducing environmental risk while increasing crop productivity and improving nutrient use efficiency in China. *J Environ Qual*. 2011;40:1051-1057. <https://doi.org/10.2134/jeq2010.0292>.
 26. Sharma HSS, Fleming C, Selby C, Rao JR, Martin T. Plant biostimulants: a review on the processing of macroalgae and use of extracts for crop management to reduce abiotic and biotic stresses. *J Appl Phycol*. 2014;26:465-490. <https://doi.org/10.1007/s10811-013-0101-9>.
 27. Udensi EU, Omovbude S, Nwachukwu JC. Response of Cucumber (*Cucumis sativus* L.) and Weeds to Plant Spacing and Weeding Regimes in a Humid Forest Agro-Ecology of Southeastern Nigeria. 2018;22(1):1-13. Article no.IJPSS.39638
 28. Dhillon NS, Sharma P, Sharma KD, Kumar P. Effect of Plant Density and Shoot Pruning on Yield and Quality of Polyhouse Grown Cucumber. *Environment & Ecology*. 2017 Oct-Dec;35(4B):3023-3026. Website: environmentandecology.com ISSN 0970-0420
 29. Quian H. Effect of pruning and spacing on yield and quality of cucumber. *Hort. Sci*. 2000;36(2):274-278.
 30. Paulo S, Lima S, Jailton FR, Jaedson CAM, Jaeverson DAS. Plant density and fruit density of muskmelon. *Rev. Bras frutic*; c2003, 25(2). Jaboticabal.