



E-ISSN: 2663-1067

P-ISSN: 2663-1075

www.hortijournal.com

IJHFS 2024; 6(2): xx-xx

Received: xx-05-2024

Accepted: xx-06-2024

Awa Marie Coll Cissé

¹ Doctoral School Sustainable Development and Society (ED2DS), Iba Der Thiam University (UIDT) BP: A967, Thies, Senegal

² National Higher School of Agriculture (ENSA) BP 296 Km 7, Khombole Road Thies, Senegal

Cheikh Dieye

Department of Plant Biology, Cheikh Anta Diop University of Dakar, BP 5005 Dakar Fann, Senegal

Papa Madiallacké Diedhiou

UFR of Agronomic Sciences, Aquaculture and Food Technologies (UFR S2ATA) Gaston Berger University BP 234, Ngallele Road Saint-Louis, Senegal

Bassirou Mbacké

Iba Der Thiam University (UIDT) BP: A967, Thies, Senegal

Ibrahima Diedhiou

National Higher School of Agriculture (ENSA) BP 296 Km 7, Khombole Road Thies, Senegal

Corresponding Author:

Awa Marie Coll Cissé

¹ Doctoral School Sustainable Development and Society (ED2DS), Iba Der Thiam University (UIDT) BP: A967, Thies, Senegal

² National Higher School of Agriculture (ENSA) BP 296 Km 7, Khombole Road Thies, Senegal

Typology of papaya-producing farms in the Niayes area (Senegal)

Awa Marie Coll Cissé, Cheikh Dieye, Papa Madiallacké Diedhiou, Bassirou Mbacké and Ibrahima Diedhiou

DOI: <https://doi.org/10.33545/26631067.2024.v6.i2a.228>

Abstract

Senegal's agricultural landscape is marked by the growing importance of papaya cultivation, which is increasingly found on farms. Papaya cultivation is emerging to make investments profitable, particularly in the Niayes area. Despite its importance, the papaya sector has not been the subject of many studies. This study aims to draw up a typology of papaya-producing farms in the Niayes area. The survey results of fifty (50) producers reveal a new profile of producers: 86% have a profession other than agriculture. Only seven (7) farms have legal status. Forty-three (43) farms are less than ten (10) hectares in size. The results also show great diversity in implementing the technical itinerary and the varieties grown. Citrus growing is frequently found in association with papaya cultivation. To our knowledge, this is the first article on the subject in Senegal.

Keywords: *Carica papaya*, Senegal, technical itinerary, fertilization, varieties, cropping associations

Introduction

The Niayes area, known for its particular climatic and ecological features, is rich in potential that has earned it its agricultural vocation. It is a favored area where animal and crop production are combined in farming systems^[4].

In recent years, many agricultural businesses have been set up. The arboricultural sector in particular has seen growing interest, with the continued establishment of purely commercial farms. Agribusiness has thus become a new focus of interest for the Senegalese population, particularly those living in the Niayes. It responds to unemployment and the modernization of what is still traditional agriculture.

Horticulture has always played a key role in the Niayes area. Among these horticultural activities, fruit growing is undoubtedly a sure way of supplementing and diversifying income from vegetable growing. A wide variety of tree species are grown here, including citrus, soursop, coconut, banana, pomegranate, mango, and papaya. Year after year, citrus fruits and papaya trees take over the cultivated areas.

Recent years have seen a growing number of papaya farms in Senegal, with crops destined for export and the local market. Papaya is one of the priority crops mentioned in the Programme to Accelerate the Pace of Senegalese Agriculture (PRACAS), the agricultural component of the Emerging Senegal Plan (PSE). Paradoxically, the papaya sector suffers from a serious lack of bibliography, which prevents a mapping of the sector.

This study aims to draw up a typology of papaya-producing farms in the Niayes area. Specifically, the aim is to:

1. Identify the different types of farms involved in papaya production;
2. review the technical itineraries used.

Materials and methods

Study site

The surveys were carried out in fifty (50) farms located in the Niayes area, precisely in Peyckouck, Fandène, Notto Diobass, Sindia, Pout, Bayakh, Diender, Bambilor, Potou, Mboro, Notto Gouy Diama and Diogo (Fig. 1).




Fig 1: Location of farms visited across the Niayes

Survey and investigation

An extensive survey was carried out to sample and characterize the papaya plantations. We used the snowball method, which consists of distributing the questionnaire to people with the characteristics we are looking for and then asking them to refer us to others with a similar profile for the rest of the survey. The snowball method is chosen because of the lack of a database.

The growers' contacts are obtained from local professionals helping them to introduce papaya to their farms, or from nurserymen. After each survey, the contacts of neighboring growers were collected. Fifty (50) farms were visited.

Data was collected using observations and surveys focusing mainly on producers, the varieties chosen, previous crops, and cropping techniques used: cropping calendar, irrigation, fertilization, crop rotation, crop associations, harvesting techniques, packaging, and yield. Data on the geolocation of the farms was also recorded.

Data analysis

The data were recorded and analyzed using Excel spreadsheets. The maps were drawn using QGIS software.

Results

Profile of producers

The survey showed that 96% of growers were male, compared with 4% who were female (Fig. 2).




Fig 2: Breakdown of producers by gender

Most of the growers we met were young men. This corroborates the strong male dominance of Senegal's agricultural sector, which is linked to the difficulty women have in gaining access to land and to the traditional reference to land management, which they usually acquire indirectly through inheritance [2, 3]. This is also explained by the fact that women are mainly involved in selling fruit.

The average age is 44, with a minimum of 28 and a maximum of 70. Most of the farmers are educated (66%), with 12% having at least a middle school leaving certificate (BFEM) and 54% having at least a baccalaureate (Fig. 3).




Fig 3: The level of education of the producers met

The strong presence of young people indicates a growing interest in agricultural entrepreneurship, which has been advocated in recent years as an alternative to unemployment and a means of retraining. The fairly advanced level of education of producers differs from the results obtained by [1] who mention that a percentage of 52.9% of producers are not formally educated. This can be explained by a reconsideration of the agricultural sector.

Producers who have a profession other than engineering, agricultural technician or traditional farming represent 86% (Fig. 4).




Fig 4: Representativeness of agricultural professionals

Agricultural professionals are in the minority among the producers we met, which shows a return to agriculture but also suggests a low level of mastery of technical itineraries due to a lack of training. Some of them are in contact with agricultural professionals for guidance, with those who sold them the nurseries or encouraged them to join the industry. This lack of expertise explains the wide variety of methods used, with everyone going their way and following the advice they receive.

Farmers' experience in papaya cultivation is fairly recent, generally less than three (3) years. The average is 2.42 years, with a minimum of one (1) year and a maximum of seven (7) years. The low number of years of experience of papaya growers confirms the recent introduction of papaya into cropping systems. This differs from the results obtained by [1], who suggest that 52% of producers in the Niayes zone have more than fifteen (15) years of experience in market gardening.

Review of farming practices

Farm organisation

Surface area

The farms visited vary in size from 0.03 to 53 ha with an average of 5.28 ha. The majority of orchards, i.e. forty-three (43) have an area of less than 10 ha (Fig. 5).




Fig 5: Distribution of farms by total area

The distribution of farms according to area shows the presence of large landlords with a clear interest in papaya, which is a high-potential crop, and corporate farms with areas of several dozen hectares, which is in line with the results of [8]. This distribution also shows the presence of small producers who see papaya as a profitable crop and hope to increase the productivity of their small areas.

A difference was noted in the distribution of land according to the agricultural activities carried out on it. Thus part of the farm is allocated to papaya or the entire area. Papaya trees are most often planted between the rows when there is a cropping association. The plants may be placed in separate plots depending on the planting period or spacing used.

The percentages of area allocated to papaya range from 1 to 100% of the plot.

Labour and farm management

Only seven (7) of the fifty (50) farms have legal status. There are two (2) Simplified Joint Stock Companies (SAS), three (3) Economic Interest Groups (GIE), one (1) Sole Proprietorship, and one (1) One-person Limited Liability Company (SUARL). It should be noted that the low level of formalization of the activity is felt in the absence of registers to record the various treatments or even the exact quantities harvested.

Two (2) types of management are noted: the owner himself is the manager or he delegates the organization of production. In 33 out of 50 cases, the owner hires workers but manages the production activities himself. In the remaining cases, the owner employs either a horticultural technician, an agricultural engineer, or an experienced worker.

The average number of workers per farm is 3.46, with a minimum of one (1) worker and a maximum of twenty (20) workers. The ratio per hectare is a minimum of 0.5 and a maximum of 33.33 for land used for housing that has been transformed into an orchard and is therefore very small.

The farms visited were not family farms. This differs from the results obtained by [6]. These farms are set up for purely entrepreneurial reasons, some of which are associated with a

passion for agriculture or fruit growing in particular. The workforce is not family-based.

Criteria for choosing papaya

The growers surveyed all mentioned the rapid return on papaya cultivation. In fact, despite the heavy investment required, there are no marketing constraints. Demand far outstrips supply. Banas-banas (wholesale traders) sometimes place orders before the harvest or come to the field to buy. Papaya stands out as a noble fruit that does not suffer from poor sales.

As well as selling well, many growers feel that papaya sells at a good price. Only two (2) growers complained about the low selling price, especially when the mango is available on the market: 300frs per kilogram. It should therefore be emphasised that it is easy to make a profit even without perfect mastery of the technical itinerary. Many were attracted to papaya after a test phase of around ten (10) plants.

Papaya is preferred to mango because of its short cycle. On several farms, it is increasingly replacing mango. Compared

with market gardening, the growers highlighted the fact that the harvest is spread out over time and that there are no financial costs associated with it (harvesting, sacking, transport). Once established, the papaya plants are only replaced after two (2) or three (3) years, whereas after less than one (1) year the market garden plants are at the end of their cycle. The land therefore has to be cleaned and restored.

Nevertheless, the lack of a bibliography in line with cultural conditions in Senegal is felt and emphasized by several growers.

Varieties grown

Seventeen (17) varieties were listed. These are : *Solo*, *Horizon*, *Red Lady*, *Red Army*, *Red Royal*, *Cortex*, *Vega*, *Linar*, *Red Nemessis*, *F1 Maradona*, *Rana*, *Red King*, *Red Queen*, *Formusa*, *Amazone*, *Red Maradol*, *Red Matador*. The varieties *Horizon*, and *Solo* were present respectively in 23, and 21 orchards. The varieties *Red Lady* and *Red Nemessis* were present in 17 out of 50 farms (Fig. 6).




Fig 6: Representativeness of the different papaya varieties

The selection criteria stated by the growers were ease of sale, fruit size, sugar content, disease resistance, productivity, shelf life, early entry into production, appearance of the fruit, and ease of harvesting. In addition, it was noted that growers paid particular attention to the advice of professionals in the field when making their choice.

Often two (2) or more varieties are combined, to diversify the offer or just to test their qualities.

There is a great deal of variability between the different varieties in terms of flesh color, size, and taste. This is in line with [5]. Dwarf varieties are the most common on farms. They are appreciated because of their small size, which makes them easier to harvest and apply phytosanitary treatments. Varieties with large fruit sizes are more difficult to sell on certain markets, but they do save a few kilos when they are weighed. Growers can therefore see how to

combine different varieties which, when marketed at the same time, can be sold easily, while making the most of the weight aspect. This explains the wide range of varieties found in orchards. It should be pointed out that there is a mix-up between varieties with similar characteristics to Solo and those that are not. On the market, medium-sized red-fleshed varieties are mostly referred to as Solo by merchants or buyers. The latter is the most widely grown variety in the literature [7, 9], which differs from the results obtained and can be explained by the emergence of new, better-performing varieties and the development of the sector.

The planting periods are also very varied and can be explained either by the fear of pest pressure during the rainy season, or by the calculation that the harvest will coincide with the scarcity of mango, and therefore the sale of papaya at a better price.

Cultural associations

Citrus trees, particularly lemons, are strongly represented on the farms (44/50). Next come mango and soursop, found on

17 and 14 farms respectively. Other associations are noted with pomegranate, lemon, orange, coconut, pomelo and banana. These are located in the inter-rows (Fig. 7).




Fig 7: Representation of different arboreal species

Citrus fruits have always been well represented in the Niayes, previously in association with mango, as shown by the work of [8]. Papaya is increasingly taking the place of mango in orchards, taking into account the results obtained. There is an association with market garden species. These include tomatoes, parsley, chilies, zucchini, watermelons, onions, peppers and eggplants. They are either planted in

rows to save space or on separate plots. The large number of species (16) found on farms confirms the area's market gardening vocation. Some farmers raise laying hens, broilers, guinea fowl, ostriches, fattening cattle, sheep and goats, dairy cows, and fish (Fig. 8).




Fig 8: Different agricultural activities found on farms

Fig. 9 shows the representativeness of the different animal production activities on the farms. We will group all the poultry we found under the heading of chicken coop. This is

the most common activity, as it was found on twenty (20) farms.




Fig 9: Different livestock production activities

There is a great diversity of farming activities, which can be explained by the search for sources of income that can support the farm and family expenses. This confirms the findings of [8].

Spacing

A wide range of spacings was noted, sometimes on the same farm but usually for testing purposes: 1 m X 1 m, 1,5 m X 2 m, 1,5 m X 1m, 2 m X 2 m, 2,5 m X 2 m, 2,5 m X 2,5 m, 2,5 m X 10 m, 3 m X 3 m. High density is preferred by some growers to make the most of or increase the productivity of a small area to their liking. Others avoid high densities because of pest pressure and poor sunlight.

Another group set up very wide spacings to place other fruit trees in the spaces between the rows. Nevertheless, the 2 x 2 m spacing is the most common, with 22 farms out of 50.

Fertilisation

Bottom dressing

The use of manure during harvesting is a common point between producers. This manure comes from a variety of sources: poultry, beef, sheep, donkey, horse, and goat. It is often collected in the orchard from the hen houses or paddocks. Beef and poultry manure are the most frequently used (Fig. 10).




Fig 10: Type of manure

Some farmers add DAP (Diammonium phosphate) to the manure during the season. The quantities added are difficult to quantify. They are of the order of a few shovels or buckets for manure and teaspoons or matchboxes for DAP.

Maintenance manure

A variety of mineral fertilizers are used for maintenance: 10-10-20, 13-13-13, 15-15-15, urea, DAP, magnesium nitrate, potassium nitrate, potassium sulfate, MAP (Monoammonium phosphate), calcium, potassium chloride,

18-18-18, 10-20-20, 20-5-5, humic acid. Urea is used at the start of cultivation on young plants. Only a small number of growers continue to apply it throughout the cycle. Potassium nitrate and potassium sulfate are used during the fruiting phase to increase the sugar content of the fruit. The other fertilizers mentioned above are applied during the cycle. Quantities are difficult to measure and correspond to either a teaspoon, a matchbox, or a relatively small handful, depending on the size of the plant. Urea is the most commonly used mineral fertilizer, followed by 10-10-20 and

potassium sulfate (Fig. 11).




Fig 11: Mineral fertilizers used by growers

Soluble fertilizers are fertigated on some farms. These include the following fertilizers: MAP, 18-18-18, urea, humic acid, potassium chloride, and potassium sulfate. Fertigation is used by 16% of growers.

Foliar fertilizers were used by sixteen (16) of the fifty (50) growers interviewed. Those found in the field are LOBI 44, FOLIGA, GROFOL 20-30-10, 8-8-40, 10-50-6, 17-17-17, 4-16-28, SYNERGIAQUE, FORCLOP GOLDEN 4-14-4. They are often sprayed at the same time as crop protection products.

On fish farms, irrigation is sometimes carried out from the rearing tanks, taking advantage of the minerals dissolved in the fish droppings.

Fertilization schedules are set at a moment's notice: when the manure in the trough is exhausted, for example, depending on the sugar level in the fruit, on any growth delays noted, or on the observation of weakened plants following a pathogen infestation. For others, fertilizers are applied systematically every month, every two (2) months, or according to the different phases of growth, in this case, flowering and fruiting.

The aforementioned poor mastery of technical itineraries explains the problems encountered in the dosage and choice of fertilizers. The interest on the part of stakeholders in finding a way to increase yields also justifies the non-moderate use of mineral fertilizers observed in the field. The effectiveness of the products is conveyed by word of mouth and explains the similarity noted between the products used in neighboring orchards. Of the fertilizers recommended by RADHORT (African Network for Horticultural Development), only urea, potassium sulfate, and manure are used by growers [7].

Drainage works and irrigation

The drainage works encountered are wells fitted with pumps and boreholes/mini-boreholes. The installation of these structures is accompanied by that of basins or water towers. Some farms use both types of water sources. The energy

source is either electric with a fuel-powered generator, or more frequently solar energy. Some farmers have access to water from nearby community boreholes (Fig. 12).




Fig 12: Source of irrigation water for papaya farms

The large number of wells and boreholes indicates an outcrop of the water table and therefore extensive exploitation of groundwater [4, 8].

It is difficult to quantify the quantities of water used. Irrigation frequency is from daily to one (1) day in 4 for adult plants. It is daily for young plants. The frequency with which the plants are watered depends on the availability of water, the area to be irrigated, and the type of soil. Irrigation is daily on predominantly sandy soils.

Irrigation is carried out using a drip system, hosepipes, sprinklers, or watering cans. The drip system is the most common with 48%, followed by hoses with 44%. Then come sprinklers (6%) and watering cans (2%).

Harvesting and yield

Harvesting begins between three (3) and nine (9) months depending on the variety. In the event of an attack, the start of production is extended. Thirty (30) farms harvest between six (6) and nine (9) months. All the cut fruit will

already have started to turn, and harvesting is done by hand without the use of tools for dwarf varieties. A pole is used for the classic varieties.

The interval between harvests varies according to the season. In hot weather, harvests are closer together and take place almost every week. In cooler periods, the interval between two (2) harvests can vary from ten (10) to fifteen (15) days. Estimating the harvest remains difficult. For some, quantities are calculated approximately per plant/week, for others per plant/10 or 15 days, and for others per harvest over the entire area, with no fixed intervals. Others stop counting when they have a return on their investment.

In addition, the major factor highlighted by the growers is the high productivity of the plants, whatever the variety and the level of mastery of the technical itinerary. During the cruising phase, yields vary from ten (10) kg per week to six (6) tonnes per week, with a positive correlation with the number of plants.

Few growers are interested in processing. Only three (3) producers are processing (dried papaya or juice). Papaya is a fruit that can be sold easily and at a good price for producers. It does not suffer from poor sales, but rather from shortages due to high demand. Given this situation, growers do not feel the need to set up storage facilities or process the fruit. This justifies the small number of growers involved in processing.

Conclusion

This study reveals a wide range of cultivation practices for papaya in the Niayes area, partly due to a lack of reference literature. It shows a new type of Senegalese farmer, with educated producers open to agribusiness. The papaya sector is seen in the Niayes area as an opportunity to be seized by the Senegalese horticultural sector.

References

1. Boinahadji AK, Kébé B. Rapport d'enquête sur la reconnaissance et la gestion des insectes ravageurs des cultures maraîchères par les producteurs de la zone des Niayes. *Eclasio*; c2018. p. 23.
2. Diop A. Diagnostic des pratiques d'utilisation et quantification des pesticides dans la zone des Niayes de Dakar (Sénégal). France: Université du Littoral Côte d'Opale; c2013.
3. Diop F, Touré L. Contribution du système d'information géographique (SIG) dans la lecture du problème de l'accès des femmes au foncier rural au Sénégal : cas des communautés rurales de Keur Momar Sarr, Médina Ndiathbé (Vallée et Zone sylvo-pastorale), Diender (Niayes), Bandafassi (Sénégal Oriental), Diendé (Casamance). *Géographie du Laboratoire Leïdi*. 2012;10:18.
4. Fall ST, Fall AS. Cités horticoles en sursis - L'agriculture urbaine dans les grandes Niayes au Sénégal. Ottawa (Ontario), Canada: Centre de recherches pour le développement international; c2001. p. 150.
5. Koffi NBC. Pourriture de racines et/ou du collet du papayer (*Carica papaya* Linn. Var. golden) en Côte d'Ivoire: caractérisation de l'agent pathogène, évaluation des traitements chimiques et perspectives de lutte biologique. Côte d'Ivoire: Nangui Abrogoua; c2013.

6. Ndaw N, Sow SA, Faye CAT, Diallo ML. Activités extractives minières et horticulture dans les Niayes de la Commune de Darou Khoudoss, Sénégal. *Afrique SCIENCE*. 2019;15(6):130-42.
7. Telemans B. La culture du papayer au Sénégal. Dakar: Institut Sénégalais de Recherche Agricole (ISRA); c2012. p. 14.
8. Touré O, Seck SM. Exploitations familiales et entreprises agricoles dans la zone des Niayes au Sénégal. International Institute for Environment and Development; c2005.
9. Yogiraj V, Goyal PK, Chauhan CS, Vyas B. *Carica papaya* Linn: an overview. *Int. J Herb Med*. 2014;2(5):1-8.