



E-ISSN: 2663-1067
P-ISSN: 2663-1075
IJHFS 2020; 2(2): 05-09
Received: 04-06-2020
Accepted: 06-07-2020

Eleduma AF
Department of Agricultural
Technology, Rufus Giwa
Polytechnic, P. M. B. 1019,
Owo, Ondo, Nigeria

Aderibigbe ATB
Department of Crop, Soil and
Pest Management Technology,
Rufus Giwa Polytechnic,
P. M. B. 1019, Owo, Ondo,
Nigeria

Sanni KO
Department of Crop
Production and Horticulture,
Lagos State Polytechnic,
Ikorodu, Lagos, Nigeria

Obabire SO
Department of Crop, Soil and
Pest Management Technology,
Rufus Giwa Polytechnic,
P. M. B. 1019, Owo, Ondo,
Nigeria

Corresponding Author:
Sanni KO
Department of Crop
Production and Horticulture,
Lagos State Polytechnic,
Ikorodu, Lagos, Nigeria

Effect of cattle manure on soil nutrients dynamics and performances of maize (*Zea mays* L) grown on sandy loam soil in forest-savannah transition zone southwest Nigeria

Eleduma AF, Aderibigbe ATB, Sanni KO and Obabire SO

Abstract

The contribution of inorganic fertilizer to enhance crop growth and yield cannot be ignored, but on the other hand their indiscriminate use is causing deterioration of the soil structure and soil acidity. A field experiment was conducted at the Teaching and Research Farm of Rufus Giwa Polytechnic, Owo, during the wet season of 2015 to study the effect of variable rate of cattle manure on the growth and yield of maize (*Zea Mays* L). Five rates of cattle manure: control, 5, 10, 15, and 20 t/ha⁻¹ replicates three times laid out in a Randomized Complete Block Design (RCBD). Data were collected on vegetative traits and yield component of the maize and statistically analyzed. The growth and yield attributes measured in the field include number of leaves, plant height (cm), leaves length (cm), leaves area (cm), stem girth (mm), number of cobs, cobs length (cm), cobs circumference (cm) number of grain per cobs, cobs weight (kg), grain weight (kg), and husk weight (kg). The growth and yield attributes increased significantly accordingly to the rates of cattle manure due to continuous supply of nutrient. The control plot gives significantly lower means and maize that received 20 t ha⁻¹ recorded the highest mean in all growth and yield parameter evaluated in the study. In conclusion, applications of cattle manure at the rate of 20 t ha⁻¹ significantly improve vegetative growth and yield attributes of maize plant and therefore recommended that resource-poor farmer can adopt the use of cattle manure at 20 t ha⁻¹ as substitute for inorganic fertilizer, thus reduce environmental pollution posed by the disposal of the cattle waste and ultimately improve yield in the study area.

Keywords: cattle manure; substitute; vegetative growth; yield attributes

Introduction

Maize, sometimes referred to as corn is ranked with wheat and rice as one of the most produced and consumed cereal crop in Nigeria (Vaidya, 1986) [39]. In spite of its nutritional and economic value its production has been hampered by low yield caused by several factors and chief among it is low soil fertility status. Nigeria soil has a high potential for crop production yield, levels obtained under farmer's condition are usually low due to poor soil management and conservation method. The type of problem is solved through the use of either inorganic or organic fertilizer. However, the use of inorganic fertilizer by resource poor farmers is limited by its scarcity and cost (Akanbi *et al.*, 2001) [7] and untimely availability (Adedoyin, 1995) [5]. Under intensive agriculture, the use of inorganic fertilizer has not been helpful because it is associated with soil acidity, nutrient imbalance and reduction in crop yield (Kang and Juo, 1980; Obi and Ebo, 1995) [23, 25]. The use of organic manure to meet the nutrient requirement of crop would be an inevitable practice since organic manures generally improve the soil physical, chemical and biological properties along with conserving capacity of the moisture holding capacity of soil and this, resulting in enhanced crop productivity along with maintaining the quality of crop produce (Mahhesarappa *et al.* 1999) [24]. Although organic manures contain plant nutrients in small quantities compared to inorganic fertilizers, the presence of growth promoting principle like enzymes and hormones, besides plant nutrient make them essential for improvement of soil fertility and productivity (Bhurna, 2001) [13]. Improvement in the public health and environmental condition are also the strong reason for advocating the use of organic material (Aboul El-majd *et al.* 2006) [3]. In the present chemical input farming system, natural (use of organic amendment *viz.* green manure, poultry manure, farmyard manure etc)

can be considered to be a solution to many problems for different cropping system in different agro-ecological zones (Sharma and Militra, 1991) [35].

With the problem associated with inorganic fertilizers already being viewed this project work was carried out to evaluate the effect of cattle manure on the growth and yield performances of maize in forest Savanah transition zone south western Nigeria.

Results

Physiochemical Properties of Soil and Cattle Manure

The physiochemical properties of the soil (0.30cm depth) at the experimental field before cropping showed that the soil textural classification was sandy loam, slightly acidic in nature with pH value of 5.49 and low and low in fertility as indicated by the low content of organic carbon, organic matter, total nitrogen, available phosphorus and exchangeable cations (Table 1).

Soils with low levels of nutrients need to be boosted with soil amendments input in form of inorganic or organic fertilizer in order to improve them for crop production. The low soil fertility status and its acidic nature are expected to benefit from application of cattle manure (Table 2).

Low soil nutrient of the experimental site was probably as a result of: (1) high proportion of sand content of the soil, which implies that basic cations would be leached more easily as texture determines the degree of retention or ease of leaching of basic cations (Wapa and Oyetayo, 2014) [40]. (2) The low NPK levels observed in the soil can be attributed to continuous cropping and increased land use intensity.

At crop maturity, application of different rate of cattle manure slightly increased the soil pH value and total nitrogen ranges from 5.58 to 5.87 and 0.12-0.28 respectively. Percentage organic carbon, Ca and Mg was significantly increased in soil amended with cattle manure compared with control plot and the highest recorded from plots supplied with 20t/ha⁻¹, while available P, Na and K was drastically and slightly decreased across the different level of application respectively (Table 1).

The marginal increase observed in the soil pH level indicates the buffering capacity of the cattle manure in reducing soil acidity. This finding however did not agree with the findings of Adekiya and Agbede (2010) [6]; Sanni (2016) [34] who observed that soil pH tended to reduce with a rise in the amount of poultry manure and application of cow manure, suggesting that poultry and cow manure led to increased acidity in the soil.

The increases in soil organic carbon, Ca and Mg observed at crop maturity from plots incorporated with varying level of cattle manure might be as a result of the slow rate in which their nutrients are released into the soil (Olowoake and Adeoye, 2013) [28] or might be due to the increased in the amount of nitrogen present in the various rates of compost applied which improved microbial activity that led to enhanced production and mineralization of organic matter from natural (native) source in soil (Sanni, 2016) [34].

The higher organic carbon content in the plot treated with cattle manure relative to other treatment plots agreed with findings of Esawy *et al.* (2009) [17] on the increase in organic matter as a result of addition of rice straw compost. While the low P, K level and marginal increase of K observed at crop maturity might be as a result of NPP uptake for the growth and development of the plant.

Table 1: Soil physiochemical properties before planting and at crop maturity

Constituents	Pre-cropping		Post-cropping		
	5t/ha ⁻¹	10t/ha ⁻¹	15t/ha ⁻¹	20t/ha ⁻¹	
Chemical					
pH (H ₂ O)	5.49	5.64	5.58	5.76	5.87
Organic carbon (%)	0.54	0.82	1.43	2.09	3.11
Total Nitrogen (%)	0.08	0.12	0.22	0.28	0.26
Available P (mg/kg)	9.18	4.91	3.82	4.32	4.81
Na (cmol/kg ⁻¹)	0.93	0.38	0.70	0.77	0.73
Ca (cmol/kg ⁻¹)	0.42	1.10	1.53	1.81	1.83
K (cmol/kg ⁻¹)	0.96	0.24	0.18	0.23	0.27
Mg (cmol/kg ⁻¹)	0.20	0.50	1.04	2.64	2.97
Physical					
Sand %	47	64.80	70.5	72.4	71.9
Silt%	44	8.0	15.7	15.0	12.4
Clay %	9.0	27.20	13.8	12.6	15.7
Textural class	Sand loam		Sandy loam		

Table 2: Chemical analysis of cattle manure used

Cattle manure	value
pH (H ₂ O)	5.48
Organic carbon (%)	2.47
Organic matter (%)	4.26
Total N (mg/kg)	1.09
Available P (mg/kg)	23.04
K (mg/kg)	1.05
Na (mg/kg)	3.48
Ca (mg/kg)	1.00
Mg (mg/kg)	0.50

Effect of Cattle Manure on mean Vegetative Growth parameters of Maize

The response of maize to the varying levels of cattle manure of the vegetative growth (plant height, number of leaves, leaves area and stem girth) is presented in Table 3. The response of plant in terms of vegetative growth followed the same trend whereby the lowest values were recorded for control plants and higher values for fertilized plants in line with the amount of manure applied i.e. the higher the amount of manure applied the greater the mean value of the vegetative growth parameters. This result suggests that nutrient availability, especially NPK, determines plant vegetative growth and development (Adebayo *et al.* 2009) [4]. The consistent poor performance of control plots (non-fertilized) and those with low level of manure revealed that when nutrients are available in adequate amounts, plant tends to grow at their optimum potential. These improvements in growth parameters with increase in rates of cattle manure applied agree with the findings of Fawuzi (1977) [18]; Uzo (1971) [38] and Aminifard, *et al.*, (2010) [8].

Table 3: Effect of Cattle Manure on mean Vegetative Growth parameters of Maize

Treatments	Plant height (cm)	Number of leaves	Leaves area (cm ²)	Stem girth
Control	102.25 ^{cd}	13.07 ^d	313.62 ^d	22.13 ^b
5 tha ⁻¹	103.15 ^c	13.18 ^c	378.07 ^c	22.94 ^b
10 tha ⁻¹	114.69 ^b	13.42 ^b	379.91 ^c	22.94 ^b
15 tha ⁻¹	118.15 ^{ab}	13.93 ^{ab}	381.136 ^b	23.43 ^b
20 tha ⁻¹	119.03 ^a	14.27 ^a	403.04 ^a	24.35 ^a
LSD (0.05)	0.81	0.06	0.31	0.86

Mean followed by the same letter (s) in a column are not significantly different according to multiple comparison, LSD $p < 0.05$.

Leaf area index (LAI) is a crucial growth index determining the capacity of plant to trap solar energy for photosynthesis and has marked effect on growth and yield of plant. The influence of cattle manure on leaf area remained significant under different application levels; maximum leaf area index (403.04) was recorded at 20 ton ha⁻¹ which was statistically higher than 15 ton ha⁻¹ (381.13), 10 ton ha⁻¹ (379.91), 5 ton ha⁻¹ (378.07) and 0 ton ha⁻¹ (313.62). However, no significant was observed among the treatments with plants amended with 10 ton ha⁻¹ (379.91) and 5 ton ha⁻¹ (378.07) cattle manure. Nitrogen (N) is one of the main plant nutrients affecting plant growth and yield (Tafteh and Sepaskhah, 2012) [37] and leaf area increase with increase in N level (Oscar and Tollennar, 2006) [30]. Increased leaf area in soil amended with organic fertilizer could probably be attributed to N availability which promoted leaf area during vegetative development and also helped to maintain functional leaf area during the growth period (Cox *et al.* 1993) [14].

Stem diameter was significantly ($p < 0.05$) influenced by varying level of cattle manure application. Application of 20 t ha⁻¹ cattle manure gave the highest response in respect, which was followed by application of 20 t ha⁻¹, while the control treatment gave the least plant diameter. Mean treatment value obtained from plant amended with 15, 10, 5 and controls were similar and in comparison were not significantly different. Organic manures have been said to improve soil fertility by activating soil microbial biomass, which in turn leads to development in corps (Ayuso *et al.*,

1996) [12] and this may have been responsible for the observed increase in stem diameter resulting from nutrient application. Plant diameter would have positive implication on lodging, particularly during fruiting; the thicker the stem, the less likely the plant would lodge as a result of fruit carriage or other lodge inducing factors, such as wind.

Plants provided with 20 t ha⁻¹ of cattle manure were significantly ($p < 0.05$) taller than the other levels of manure and control plants. The steady increase in height of maize plants with increase in rate of cattle manure suggests that quantity of manure applied affects nutrients availability for uptake by plants which promoted vigorous plant growth through efficient photosynthesis (Ayoub *et al.* 1991; Islam, 2002; Iqtidar *et al.* 2006) [11, 21, 20].

The highest number of plant leaves in the present study was recorded in plants amended with 10 kg m⁻² of organic fertilizer which was significantly ($p < 0.05$) higher than the control but not statistical different with plants amended with 5 and 7.5 kg m⁻². The application of cattle manure to the soil at varying levels may have improved soil fertility and soil structure, increased soil organic matter and enhanced microbial activity (Rehman *et al.* 2010) [33] and the nutrients released from manure thus support and enhanced rapid root development (Abou El-Magd *et al.* 2005) [2] which might have enhanced leaf growth.

Effect of cattle manure application on mean yield attributes of maize

Table 4: Effect of cattle manure application on mean yield attributes of maize.

Treatments	No of Cobs	No of Grains/ Cob	Cob Length (cm)	Cob Diameter(cm)	Grain Weight (Kg)	Husk Weight (Kg)
Control	5.33 ^a	400.12 ^{de}	15.05 ^c	3.49 ^c	1.13 ^c	0.11 ^a
5 tha ⁻¹	5.33 ^a	439.67 ^d	15.43 ^c	3.77 ^c	1.24 ^c	0.16 ^a
10 tha ⁻¹	5.67 ^a	447.40 ^c	15.69 ^c	3.66 ^c	1.54 ^c	0.15 ^a
15 tha ⁻¹	5.67 ^a	473.49 ^b	16.61 ^b	3.94 ^c	1.70 ^b	0.20 ^a
20 tha ⁻¹	6.07 ^a	516.83 ^a	17.38 ^a	4.18 ^c	2.04 ^a	0.21 ^a
LSD (0.05)	0.89	0.41	0.41	0.63	0.39	0.81

Mean followed by the same letter (s) in a column are not significantly different according to multiple comparison, LSD $p < 0.05$.

A non-significant treatment effect was revealed from mean value showed in Table 4 for number of harvested cob per plant, cob diameter and husk weight, however maize plants amended with 20 t ha⁻¹ had superior absolute treatment mean numbers which decreased with each decrease in level of cattle manure applied. Meanwhile, the treatment mean obtained for number of grains per cobs, cob length and grain weight increased accordingly to the increase in the cattle manure application rates; with 20 t ha⁻¹ having the highest mean value which was significant different from mean value obtained from other application rates and control plots recorded the lowest mean treatment value. But cattle manure applied at the rates of 10 and 5 t ha⁻¹ did not differ significantly compared to control treatment. The highest number of grains, cob length, grain weight and husk weight obtained from 20 t ha⁻¹ was as a result of higher nutrient base supplied by the rate when compared with other rates where there was reduction with decrease in rate of cattle manure applied. The higher yield attributes obtained from the application of higher rates of cattle manure is in accordance with the studies of Devi *et al.*, (2002) [15]; Ogar and Asiegbu, (2005) [26]; Aujla *et al.*, (2007) [10]; and Aminifard, *et al.*, (2010) [8] where higher rates of nutrients

increased the average crop weight and volume. This might be attributed to the stimulating effect of cattle manure 480 that supplies plant with nutrients required for better yield (Abdelrazzag, 2002) [1].

The application of increasing levels of cattle manure increased the maize growth, which might have been due to the balance availability of nutrients to the plants that resulted in a favourable soil environment. These favourable conditions increased the nutrient availability and water holding capacity of the soil resulting in enhanced growth and yield (Rashid *et al.*, 2013) [32]. Therefore, it is recommended that resource-poor farmers in the study area should utilize cattle manure at the rate of 20 t ha⁻¹ for sustainable cultivation and optimum yield of maize in the study area as the manure contribute to the improvement of the low fertility status of the soil.

Materials and Methods

Experimental layout and land preparation

The field experimental was laid out in a randomized complete block design (RCBD) replicated three (3) times with five treatments on a 104m² area of land at the Teaching, Research and Commercial farm, Rufus Giwa

Polytechnic, Owo, Ondo state, Nigeria, located on latitude 7° 12'N, longitude 5° 35'E within the forest savanna transition zone, south western Nigeria.

The land was occupied by *Sida acuta*, *Imperata cylindrica*, Siamweeds and some other common weeds, the land had been under continuous cultivation for the past four years without application of either inorganic and organic fertilizers. The land was cleared and stumps manually and thereafter ploughed and harrowed mechanically in order to obtain a clean fine hit soil. The experimental land was divided into 3 blocks of 12m x 2m (24 m²) each with a distance of 0.50m between each plots and each plot size was 2m x 2m (4m²) which produced a total number 15 plot. The treatments are: T₁ – 0tons/ha of cattle manure, T₂ – 5tons/ha of cattle manure, T₃–10tons/ha of cattle manure, T₄ – 15tons/ha of cattle manure and T₅ – 20tons/ha of cattle manure. The cattle manure was applied to the plot 2 week after planting (WAP) using ring method of application after being pulverized.

Crop Establishment and Management

Two seeds of improved maize seed downy middle resistance early streak resistance yellow (DMR-ESR-Y) obtained from National Center for Genetic Resource and Biotechnology (NACGRAB) Ibadan, Oyo State Nigeria was per hole at a spacing of 50cm x 50cm at a depth of 3 to 5cm; which was later thinned to one healthy plant per stand. Immediately after planting Atrazine mixed with Paraquat was applied as pre-emergence and contact weed control, when necessary rouging was carried out to eliminate emerging weeds.

Cow manure and soil analysis

Cattle manure obtained from the livestock session of the Teaching, Research and Commercial Farm, Rufus Giwa Polytechnic, Owo, Ondo State, Nigeria. The sample was air dried, crushed through a 2mm sieve and subjected to laboratory analysis following standard procedure in order to determine the chemical properties (IITA, 1989) [19]. Prior to the commencement and at the end of the experiment, top soil samples of 20 cm- 30 cm deep were collected randomly from five spots using soil auger from the experimental plot and mixed together to form a composite sample. It was air dried, sieved with a 2mm mesh-size sieve and taken to the laboratory to determine the soil's physicochemical properties using standard laboratory procedures (Olsen *et al.* 1954 [29]; Jackson, 1973 [22]; Page *et al.* 1982 [31]; Dirk and Hargarty, 1984 [16]; Okalebo *et al.* 2002) [27].

Data collection and statistical analysis

Five plants were randomly tagged from two middle rows in each plot for sampling data collection for vegetative growth and yields assessment. The following data on vegetative growth (number of leaves, leaf area, stem girth and plant height) and yield attributes (number of grains per cobs, cob circumference, grain weight and husk weight) were collected at crop maturity and harvest. Data collected from the experimental field were subjected to analysis of variance (ANOVA) using Statistical Package for Social Science (SPSS Inc., 2011) [36]. Significant treatment mean were compared using Duncan Multiple Range Test (DMRT) at 5% probability level.

References

1. Abdelrazzag A. Effect of chicken manure, sheep

- manure and inorganic fertilizer on yield and nutrients uptake by onion. Pakistan Journal of Biological Sciences 2002;5(3):266-268.
2. Abou El-Magd MM, Hoda Mohammed A, Fawzy ZF. Relationships, Growth, Yield of Broccoli with Increasing N, P or K Ratio in a Mixture of NPK Fertilizers," Annals of Agricultural Science Moshtohor 2005;43(2):791-805.
3. AbouElMagd MN, El- Bassiong M, Faziour ZF. Effect of organic manure with or without chemical fertilizer on growth and quality of some variety of Bioccoli plant. Journal of Applied Science Research 2006;2(10):791-798.
4. Adebayo AG, Akintoye HA, Aina OO, Olatunji M, Shokalu AO. Assessment of Organic amendments on growth and flower yield of Sunflower (*Helianthis annus*). In: Prceedings of Horticultural Society of Nigeria 11-16 October, 2009, Kano State, Nigeria 2009, 292-298.
5. Adedoyin SF. An assessment of the linkage in delivery of technologies on rehabilitation and management of soil: A paper presented at 3rd All Africa soil science conference, university of Ibadan 1995, 19-26.
6. Adekiya AO, Agbede TM. Growth and yield of tomato (*Lycopersicon esculentum* Mill) as influenced by poultry manure and NPK fertilizer. Emirate Journal of Food and Agriculture 2010;21(1):10-20.
7. Akanbi NB, Adediran JA, Togun AU, Sobulo RA. Effect of organic based fertilizer on growth yield and storage of Tomato. Biosci. Res. Communication 2001;20(15):10-16.
8. Aminifard MH, Aroiee HOSSEIN, Fatemi HAMIDE, Ameri A, Karimpour S. Responses of eggplant (*Solanum melongena* L.) to different rates of nitrogen under field conditions. Journal of Central European agriculture 2010;11(4):243-248.
9. Asiegbu JE, Uzo FO. Yield and yield components response of vegetable crops to farm yard manure rates in the presence of Inorganic fertilizer. Journal of Agriculture Pueterico 1984;68:243-252.
10. Aujla MS, Thind HS, Buttar GS. Fruit yield and water use efficiency of eggplant (*Solanum melongena* L.) as influenced by different quantities of nitrogen and water applied through drip and furrow irrigation. Journal of Science and Horticulture 2007;112:142-148.
11. Ayoub M, Guertin S, Lussier S, Smith DL. Timing and Level of Nitrogen Fertilizer Effects on Spring Wheat Yield in Eastern Canada. Crop Science 1994;34:748-756.
12. Ayuso MA, Pascal JA, Garcia C, Hernandez B. Evaluation of urban wastes for agricultural use. Soil Plant Nutrition 1996;42:105-111.
13. Bhuma M. Studies on impact of humic acid on sustenance of soil fertility and productivity of green grain. Msc (ag) Thesis, TNAU, Cenbatore 2001.
14. Cox WJ, Kalonge S, Cherney DJR, Reid WS. Growth Yield and Quality of Forage Maize under Different Nitrogen Management Practices Agronomy Journal 1993;85(2):341-347.
15. Devi HH, Maity TK, Paria NC, Thapa U. Response of brinjal to different sources of nitrogen. Journal of Vegetable Science 2002;29(1):45-47.
16. Dirk AT, Hargarty M. (eds.) Soil and plant analysis. Guide for Agricultural Laboratory Directors and

- Technologist in Tropical Region 1984, pp. 147.
17. Esawy M, Mahmoud I, Paul R, Nouraya A, Mohamme E. Rice Straw Composting and its Effect on Soil Properties 2009. http://findarticles.com/p/article/s/mi_7510/is_200907/ai_n39231352. Accessed 2nd May, 2017.
 18. Fawusi MOA. Influence of plant density and time of fertilizer application, growth characteristics, nutritional uptake and yield of tomato. *Scientia Horticulture* 1977;7(4):329-337.
 19. IITA. International Institute of Tropical Agriculture. Automated and semi-automated methods for soil and plant analysis. Manual series 1989, 7.
 20. Iqtidar H, Ayyaz KM, Ahmad KE. Bread Wheat Varieties as Influenced by Different Nitrogen Levels. *Journal Zhejiang University Science* 2006;7(1):70-78.
 21. Islam MR. Effects of Different Levels of Chemical and Organic Fertilizers on Growth, Yield and Protein Content of Wheat. *Journal of Biological Sciences* 2002;2(5):304-306.
 22. Jackson ML. Soil Chemical Analysis. Second edition. Prentice Hall of India Pvt. Ltd., New Delhi 1973, p 498.
 23. Kang BT, Juo ASR. Management of low activity clay soil in tropical Africa for crops production 1980, PP 129-133. In: Terry ER, KA Odoro and Scaveness (eds).
 24. Malieswarappa HP, Nansappa HV, Hedge MR, Balu SR. Influence of planting material, plant population and organic manure on yield of East India Galangal and on soil physiochemical and biological properties 1999.
 25. Obi ME, Ebo PO. The effects of organic and inorganic amendments on soil physical properties and maize production in a severely degraded sandy soil in southern Nigeria. *Bioresource Technology* 1995;51(2-3):117-123.
 26. Ogar EA, Asiegbe JE. Effects of fertilizer rates and cutting frequency on the marketable vegetables and pod yields in fluted pumpkin in southeastern, Nigeria. *Agro-Science* 2005;4(1):66-69.
 27. Okalebo JR, Gathua KW, Woomer PL. Laboratory Methods of Soil and Water Analysis: A Working Manual (2nd edn.) Nairobi, Kenya. SACRED Africa 2002, 128 pp.
 28. Olowoake AA, Adeoye GO. Influence of differently composted organic residues on the yield of maize and its residual effects on the fertility of an Alfisol in Ibadan, Nigeria. *Intl. J Agric. Env. Biotech* 2013;6(1):79-84.
 29. Olsen SR, Cole CV, Watanabe FS, Dean L. Estimation of available phosphorus in soils by extraction with sodium bicarbonate, U. S. Dept. Agron. Cire 1954, p 929.
 30. Oscar RV, Tollennar M. Effect of Genotype, Nitrogen, Plant Density and Row Spacing on the Area-Per-Leaf Profile in Maize. *Agronomy Journal* 2006;98(1):94-99.
 31. Page AL, Miller RH, Kuny DR. Methods of Soil Analysis. Part 2. 2nd ed., American Soc. Agron., Inc., Soil Sci. Soc. American Inc. Madison, Wisconsin 1982, 403-430.
 32. Rashid Z, Rashid M, Inamullah S, Rasool S, Bahar F. Effect of different levels of farmyard manure and nitrogen on the yield and nitrogen uptake by stevia (*Stevia rebaudiana* Bertoni). *African Journal of Agricultural Research* 2013;8(29):3941-3945.
 33. Rehman S, Khalil SK, Muhammad F, Rehman A, Khan AZ, Amanullah A, *et al.* Phenology, Leaf Area Index and Grain Yield of Rainfed Wheat Influenced by Organic and Inorganic Fertilizer. *Pakistan Journal of Botany* 2010;42(5):3671-3685.
 34. Sanni KO. Effect of compost, cow dung and NPK 15-15-15 fertilizer on growth and yield performance of Amaranth (*Amaranthus hybridus*). *International Journal of Advances in Scientific Research* 2016;2(03):076-082.
 35. Sharma AR, Militra BN. Effect of different rate of application of organic. *Journal of Agricultural Science (Cambridge)* 1991;177:313-318.
 36. SPSS Inc. SPSS Statistics 17.0, SPSS Inc., Chicago, IL, USA 2011.
 37. Tafteh A, Sepaskhah AR. Yield and Nitrogen Leaching in Maize Field under Different Nitrogen Rates and Partial Root Drying Irrigation. *International Journal of Plant Production* 2012;6(1):93-113.
 38. Uzo JO. Effects of nitrogen, phosphorus and potassium on the yield of tomato (*Lycopersicon esculentum* Mill) in the humid tropics. *Horticulture Resources* 1971;11:65-72.
 39. Vadya K. Amaranthus: The grain of the future. *Indian Farmer Digest* 1986;7(13):5-8.
 40. Wapa JM, Oyetola SO. Combining Effects of Nitrogen and Different Organic Manures on Major Chemical Properties of Typic Ustipsament in North-East Nigeria. *American International Journal of Biology* 2014;2(2):27-45.
 41. BR Bhatta, K Kaphle. Nutrition and reproductive underperformance of cattle in Nepal: A short review. *Int J Vet Sci Anim Husbandry* 2020;5(1):83-86.